arXiv:1609.00102v1 [gr-qc] 1 Sep 2016

Approximate Kerr-Newman-like Metric with Quadrupole

Francisco Frutos-Alfaro[1,*] and Paulo Montero-Camacho[2]

[1]School of Physics and Space Research Center of University of Costa Rica
[2]Center for Cosmology and AstroParticle Physics, The Ohio State University
[*]E-mail: *frutos@fisica.ucr.ac.cr*

September 2, 2016

Abstract

The Kerr metric is known to present issues when trying to find an interior solution. In this work we continue in our efforts to construct a more realistic exterior metric for astrophysical objects. A new approximate metric representing the spacetime of a charged, rotating and slightly-deformed body is obtained by perturbing the Kerr-Newman metric to include the mass-quadrupole and quadrupole-quadrupole orders. It has a simple form, because is Kerr-Newman-like. Its post-linear form without charge coincides with post-linear quadrupole-quadrupole metrics already found.

1 Introduction

Since Kerr proposed his metric in 1963 [1] multiple efforts have been directed to finding an interior solution of his spacetime [2, 3, 4, 5, 6, 7, 8, 9] or in generalizing the Kerr metric to a metric that allows a physical interior matching [10, 11, 12, 13]. Nonetheless, no physical interior solution exists. See [14] for a relatively recent prespective in the issues present in the Kerr metric that complicate the interior matching.

Furthermore, from some of the attempts of constructing an interior metric of the Kerr metric we see a trend for preference of an oblate spheroid instead of a sphere [15, 16]. This is a mathematical motivation indicating the value of exploring metrics of deformed objects. However, the physical motivation is much more simple, real astrophysical objects are not perfect spheres, hence allowing for small deformations in a rotating object is meaningful.

Moreover, the interest in spacetimes capable of describing charged distributions has always been high. In the early 1916's Reissner and Nordström [17, 18] found their metric, which described a static spherically symmetric charge distribution. Even though this metric does not handle rotation it has been used, for example, to do black hole lensing [19] and study Hawking radiation from a Reissner-Nordström black hole [20]. It is important to highlight that this kind of studies would benefit from a metric capable of describing charged objects but including rotation, and with the capabilities of allowing deformed objects, this last property rules out the Kerr-Newman metric [21].

In this article we continue our efforts in constructing a more realistic metric capable of representing a real astrophysical object. Here, we use a perturbative method which utilizes the Lewis metric [22] in order to find spacetimes with quadrupole moment while using the Kerr spacetime as a seed metric. Basically, our technique consists in cleverly changing the potentials of the Lewis metric while maintaining the cross term (rotational term). We have already applied this techinque and obtained other approximate metrics [23, 24, 25]. In comparison to our previous efforts this work

includes the addition of charge. Hence, our new metric is capable of representing a charged, rotating and slightly deformed massive object.
The usual question when computing new solutions of the Einstein Field Equations (EFE) is how to prove that a given metric would have physical meaning[1]. In order to confirm the physical legitimacy of a given metric one can expand it to its post-linear from and compare the result with the post-linear version of the Hartle-Thorne (HT) metric [10, 13]. Note that it is possible to find an inner solution to the HT metric [14], and hence if a metric has a similar form to the HT metric then an inner solution should exists. See [26, 27] for a more detailed discussion and some examples.
The rest of this paper is organized as follows: We start our perturbation method of the Kerr metric with the help of the Lewis potentials in section 2. In section 3, we obtained a new metric by means of our perturbative technique, this metric has rotation, quadrupole moments and charge. We checked that the metric is a solution of the EFE using a REDUCE program [28], this program is available upon request. We state our conclusions in section 4.

2 The Perturbing Method for the Kerr Metric

Here and in the following sections we will use the method developed by Frutos et al. in [23, 24, 25] to obtain a Kerr-Newman-like metric *i.e.* a spacetime capable of describing a slightly deformed rotating charged mass. We start by using the connection between the Lewis metric and the Kerr metric to obtain the Lewis potentials associated to our new metric. These potentials are later used in the perturbation method.
First of all, we start with the Lewis metric, which is given by [22]

$$ds^2 = -Vdt^2 + 2Wdtd\phi + Xd\rho^2 + Ydz^2 + Zd\phi^2, \tag{1}$$

where the chosen canonical coordinates are $x^1 = \rho$ and $x^2 = z$. The potentials V, W, Z, $X = \mathrm{e}^{\mu}$ and $Y = \mathrm{e}^{\nu}$ are functions of ρ and z with $\rho^2 = VZ + W^2$.
From [22] the transformation that leads to the Kerr metric is

$$\rho = \sqrt{\Delta}\sin\theta \qquad \text{and} \qquad z = (r - M)\cos\theta, \tag{2}$$

where $\Delta = r^2 - 2Mr + a^2 + e^2$, a is the rotational parameter and e is the electric charge.
Now, the Lewis potentials are chosen as follows

$$\begin{aligned}
V &= V_K\,\mathrm{e}^{-2\psi} = \frac{1}{\tilde{\rho}^2}[\Delta - a^2\sin^2\theta]\,\mathrm{e}^{-2\psi} \\
W &= -\frac{2Jr}{\tilde{\rho}^2}\sin^2\theta \\
X &= X_K\mathrm{e}^{2\chi} = \tilde{\rho}^2\frac{\mathrm{e}^{2\chi}}{\Delta} \\
Y &= Y_K\mathrm{e}^{2\chi} = \tilde{\rho}^2\,\mathrm{e}^{2\chi} \\
Z &= Z_K\mathrm{e}^{2\psi} = \frac{\sin^2\theta}{\tilde{\rho}^2}[(r^2 + a^2)^2 - a^2\Delta\sin^2\theta]\,\mathrm{e}^{2\psi},
\end{aligned} \tag{3}$$

where the potentials V_K, W, X_K, Y_K, Z_K are the Lewis potentials for the Kerr-Newman metric, and $\tilde{\rho}^2 = r^2 + a^2\cos^2\theta$. Also, $J = Ma$ is the angular momentum.
The cross term potential W is unaltered to preserve the following metric form

[1]Here physical meaning stands for having an interior metric.

$$
\begin{aligned}
ds^2 &= -\frac{\Delta}{\tilde{\rho}^2}[\mathrm{e}^{-\psi}dt - a\mathrm{e}^{\psi}\sin^2\theta d\phi]^2 + \frac{\sin^2\theta}{\tilde{\rho}^2}[(r^2+a^2)\mathrm{e}^{\psi}d\phi - a\mathrm{e}^{-\psi}dt]^2 \\
&+ \tilde{\rho}^2\mathrm{e}^{2\chi}\left(\frac{dr^2}{\Delta} + d\theta^2\right).
\end{aligned} \tag{4}
$$

These potentials guarantee that one gets the Kerr metric if $\psi = \chi = 0$. The function ψ and χ will be found approximately from the EFE.

3 The Approximate Kerr-Newman Metric with Quadrupole

As was stated in the previous section our problem has been reduced to finding the functions ψ and χ. Here we proceed to find such functions by solving the EFE perturbatively. Also, we discuss the limiting cases of the new metric.
The EFE are given by

$$
\begin{aligned}
G_{ij} &= R_{ij} - \frac{R}{2}g_{ij} = \kappa T_{ij}, \\
\nabla_j F^{ij} &= \frac{1}{\sqrt{-g}}\partial_j[\sqrt{-g}F^{ij}] = 0,
\end{aligned} \tag{5}
$$

where G_{ij} (i, j = 0, 1, 2, 3) are the Einstein tensor components, R_{ij} are the Ricci tensor components, R is the curvature scalar, $\kappa = 8\pi G/c^4$, $g = \det(g_{ij})$ and T_{ij} represent the energy-momentum tensor components, which are given by

$$
4\pi T_{ij} = g^{kl}F_{il}F_{jk} - \frac{1}{4}F^{ab}F_{ab}\,g_{ij}, \tag{6}
$$

where $F_{ij} = \partial_j A_i - \partial_i A_j$ are the electromagnetic tensor components, and A_i is the vector potential components.
The 1-form for the vector potential A can be written as follows

$$
\mathsf{A} = -\frac{er}{\rho^2}\left[\mathrm{e}^{-\psi}dt - a\mathrm{e}^{\psi}\sin^2\theta d\phi\right]. \tag{7}
$$

In addition, the 2-form for the electromagnetic tensor F can be obtained as follows

$$
\mathsf{F} = d\mathsf{A} = \frac{1}{2}F_{ij}dx^j \wedge dx^j. \tag{8}
$$

From (8) we can determine the energy-momentum tensor.
Let us highlight the terms that are neglected in our perturbative approach, which are

$$
\begin{aligned}
W^2\frac{\partial\psi}{\partial x^i} &\sim 0, \\
W\frac{\partial W}{\partial x^i}\frac{\partial\psi}{\partial x^i} &\sim 0, \\
W^2\frac{\partial\chi}{\partial x^i} &\sim 0, \\
W\frac{\partial W}{\partial x^i}\frac{\partial\chi}{\partial x^i} &\sim 0.
\end{aligned}
$$

Moreover, eliminating the terms corresponding to the Kerr metric into the Ricci tensor components, we get the Ricci tensor component of the appendix of [24]. Note that W plays the role of the rotation, since it is proportional to the angular momentum. However, the above expressions do not mean that factors of J^2 or a^2 vanish, what is being effectively restricted here are combinations of quadrupoles with the angular momentum or the rotation.
In order to obtain an expression for the Ricci tensor, we propose the following Ansatz

$$\begin{aligned} \psi &= \frac{q}{r^3}P_2 + \alpha\frac{Mq}{r^4}P_2, \\ \chi &= \frac{qP_2}{r^3} + \frac{Mq}{r^4}(\beta_1 + \beta_2 P_2 + \beta_3 P_2^2) + \frac{q^2}{r^6}(\beta_4 + \beta_5 P_2 + \beta_6 P_2^2 + \beta_7 P_2^3), \end{aligned} \tag{9}$$

where q represents the quadrupole parameter and P_2 is the usual Legendre polynomial, $P_2 = (3\cos^2\theta - 1)/2$. Furthermore, the α and β's are constants to be determined.
One can find the undetermined constants by substituting this Ansatz into the Ricci tensor components. We get a set of linear equations for these constants α, and β_n ($n = 1, \ldots, 7$). After solving these linear equations, the constants are found to be the same as in [27]

$$\begin{aligned} \alpha &= 3, \\ \beta_1 &= -\frac{1}{3}, \\ \beta_2 &= \beta_3 = \frac{5}{3}, \\ \beta_4 &= \frac{2}{9}, \\ \beta_5 &= -\frac{2}{3}, \\ \beta_6 &= -\frac{7}{3}, \\ \beta_7 &= \frac{25}{9}. \end{aligned} \tag{10}$$

Finally we have all the information we require to construct our Kerr-Newman-like metric. From (4), the metric components are given by

$$\begin{aligned} g_{tt} &= \frac{\mathrm{e}^{-2\psi}}{\rho^2}[a^2\sin^2\theta - \Delta], \\ g_{t\phi} &= \frac{a}{\rho^2}[\Delta - (r^2 + a^2)]\sin^2\theta = \frac{\sin^2\theta}{\rho^2}(aQ^2 - 2Jr), \\ g_{rr} &= \rho^2\frac{\mathrm{e}^{2\chi}}{\Delta}, \\ g_{\theta\theta} &= \rho^2\mathrm{e}^{2\chi}, \\ g_{\phi\phi} &= \frac{\mathrm{e}^{2\psi}}{\rho^2}[(r^2 + a^2)^2 - a^2\Delta\sin^2\theta]\sin^2\theta, \end{aligned} \tag{11}$$

We checked that (11) was valid up to order $O(aq^2,\, a^2q,\, Maq,\, Mq^2,\, M^2q,\, q^3)$ using a REDUCE program [28].
Now, we will focus in the limiting cases of the new metric. We summarized the limiting cases in table 1. First, note that if $e = 0$ in (11) one recovers the metric found by Frutos *et al.* in [24].

Absent physical property	Small physical property	Limiting metric
Charge	Quadrupole (linear)	Metric found in [24]
Charge	Quadrupole (quadratic)	Metric found in [27]
Quadrupole	-	Kerr-Newman
Quadrupole and rotation	-	Reissner-Nordström
Charge and quadrupole	-	Kerr
Charge and rotation	-	Erez-Rosen-like
Charge	Quadrupole and rotation	Metric found in [23]
Charge, quadrupole and rotation	-	Schwarzschild

Table 1: Limiting cases.

Therefore, following [24] the other interesting limiting cases are: The Kerr metric if $e = q = 0$, the metric found in [23] if $e = a^2 = q^2 = 0$, the Erez-Rosen-like metric described in [24] if $a = e = 0$, and the Schwarzchild metric if $a = e = q = 0$. Furthermore, one obtains the Kerr-Newman geometry if $q = 0$. Also, the Reissner-Nordström metric is obtained if $a = q = 0$. Thus, all the expected limiting cases can be obtained from this new metric.
Here we will not show the matching of (11) with the HT metric. This matching is vital because it guarantees that an interior solution exists. We will not show it since it should be trivial to follow [24] since our metric has the same form, or equivalently follow [26]. Moreover, in [27], it was shown that the multipole structure of this metric without charge is non-isometric with the Quevedo-Mashhoon [12, 13] and the Manko-Novikov [29] metrics. Then, this new metric should not be isometric with charged version of these metrics.

4 Conclusion

A metric with charge, deformations and rotation was obtained by solving the EFE pertrubatively. The expected limiting cases of this new metric were explored and found, which is a positive sign. Moreover, these limiting cases confirm that our metric adequately describes a mass with charge and quadrupole under rotation. We successfully applied the perturbation method developed by Frutos et al in [23, 24, 25, 27] to obtain a new approximate metric. Notice that the main improvement of our work with respect to [24, 27] is the inclusion of charge. Ideally, this moves us closer to representing an actual astrophysical object. Further, one could expect that careful understanding of this procedure might eventually lead us towards a magnetized object.
Even though we did not explicitly showed that the new metric can be matched to the Hartle-Thorne metric the previous success in the the non-charged metrics [24, 27] is encouraging. Particularly, because of the similarity with that non-charged metric we do not expect any issues in the matching. This matching is crucial for the sake of showing that an interior metric for our new metric exists.
Since this new metric can describe real charged astrophysical objects in a more realistic fashion than the Kerr-Newman or Reissner-Nordström metrics, then our metric should be attractive for astrophysical applications, like gravitational lensing and relativistic magnetohydrodynamics. Furthermore, computational implementation of this metric should not imply additional difficulties since it mantains a similar form to the Kerr metric.

References

[1] R. P. Kerr. Gravitational field of a spinning mass as an example of algebraically special metrics.

Physical Review Letters, 11:237–238, 1963. http://dx.doi.org/10.1103/PhysRevLett.11.237

[2] J. E. Cuchí, A. Molina and E. Ruiz Double Shell Stars as Source of the Kerr Metric in the CMMR Approximation. Journal of Physics: Conference Series, 314, (2011), Article ID: 012070 http://iopscience.iop.org/1742-6596/314/1/012070

[3] S. Haggag and J. Marek. A nearly-perfect-fluid source for the Kerr metric. Il Nuovo Cimento B, 62: 273–282, 1981. http://dx.doi.org/10.1007/BF02721277

[4] S. Haggag. A fluid source for the Kerr metric. Il Nuovo Cimento B, 105: 365-370, 1990. http://dx.doi.org/10.1007/BF02728818

[5] S. Haggag. A static axisymmetric anisotropic fluid solution in general relativity. Astrophysics and Space Science, 173: 47-51, 1990. http://dx.doi.org/10.1007/BF00642561

[6] W. Hernández. Material Sources for the Kerr Metric. Physical Review, 159, 1070–1072, 1967. http://dx.doi.org/10.1103/PhysRev.159.1070

[7] A. Krasinski. A newtonian model of the source of the Kerr metric. Physics Letters, 80A: 238-242, 1980. http://dx.doi.org/10.1016/0375-9601(80)90010-9

[8] J. P. Krisch and E. N. Glass. Counter-Rotating Kerr Manifolds Separated by a Fluid Shell. Classical and Quantum Gravity, 26, 2009. http://dx.doi.org/10.1088/0264-9381/26/17/175010

[9] A. Ramadan. Fluid sources of the Kerr metric. Il Nuovo Cimento, 119B: 123-129, 2004.

[10] J. B. Hartle and K. S. Thorne. Slowly rotating relativistic stars. II. Models for neutron stars and supermassive stars. Astrophysical Journal, 153:807–834, 1968. http://dx.doi.org/10.1086/149707

[11] K. S. Thorne. Multipole expansions of gravitational radiation. Reviews on Modern Physics, 52(2):299–340, 1980. http://dx.doi.org/10.1103/RevModPhys.52.299

[12] H. Quevedo and B. Mashhoon. Generalization of Kerr spacetime. Physical Review, 43(12):3902–3906, 1991. http://dx.doi.org/10.1103/PhysRevD.43.3902

[13] H. Quevedo. Exterior and interior metrics with quadrupole moment. General Relativity and Gravitation, 43(4): 1141–1152, 2011. http://dx.doi.org/10.1007/s10714-010-0940-5

[14] K. Boshkayev, H. Quevedo and R. Ruffini. Gravitational field of compact objects in general relativity. Physical Review D, 86, 064043, 2012.

[15] S. Viaggiu. Interior Kerr solutions with the Newman-Janis algorithm starting with physically reasonable space-times. International Journal of Modern Physics D, 15: 1441–1453, 2006. http://dx.doi.org/10.1142/S0218271806009169

[16] S. P. Drake and R. Turolla. The application of the Newman-Janis algorithm in obtaining interior solutions of the Kerr metric. Classical and Quantum Gravity, 14: 1883–1897, 1997. http://dx.doi.org/10.1088/0264-9381/14/7/021

[17] G. Nordström. On the energy of the gravitation field in Einstein's theory. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, 20(2): 1238–1245, 1918.

[18] H. Reissner. Über die Eigengravitation des elektrischen Feldes nach der Einsteinschen Theorie. Annalen der Physik, 355(9): 106–112, 1916. http://dx.doi.org/10.1002/andp.19163550905

[19] E. F. Eiroa, G. E. Romero and D. F. Torres. Reissner-Nordström black hole lensing. Physical Review D, 66: 024010, 2002. http://dx.doi.org/10.1103/PhysRevD.66.024010

[20] J. Zhang and Z. Zhao. Hawking radiation of charged particles via tunneling from the Reissner-Nordström black hole. Journal of High Energy Physics, 10: 055, 2005. http://iopscience.iop.org/1126-6708/2005/10/055

[21] E. T. Newman, E. Couch, K. Chinapared, A. Exton, A. Prakash and R. Torrence. Metric of a rotating charged mass. Journal of Mathematical Physics, 6, 918, 1965. http://dx.doi.org/10.1063/1.1704351

[22] M. Carmeli. *Classical Fields: General Relativity and Gauge Theory.* World Scientific Publishing, Singapore, 2001. http://www.worldscientific.com/worldscibooks/10.1142/4843

[23] F. Frutos-Alfaro, E. Retana-Montenegro, I. Cordero-García, and J. Bonatti-González. Metric of a Slow Rotating Body with Quadrupole Moment from the Erez-Rosen Metric. International Journal of Astronomy and Astrophysics, 3, 431–437, 2013. http://dx.doi.org/10.4236/ijaa.2013.34051

[24] F. Frutos-Alfaro, P. Montero-Camacho, M. Araya, and J. Bonatti-González Approximate Metric for a Rotating Deformed Mass. International Journal of Astronomy and Astrophysics, 5, 1–10, 2015. http://dx.doi.org/10.4236/ijaa.2015.51001

[25] P. Montero-Camacho, F. Frutos-Alfaro, C. Gutiérrez-Chaves. Slowly rotating Curzon-Chazy Metric., Revista de Matemática (Teoría y Aplicaciones) 22 (2):265–274, 2015. http://dx.doi.org/10.15517/rmta.v22i2.20833 (http://arxiv.org/abs/1405.2899)

[26] F. Frutos-Alfaro and M. Soffel. On the Post-linear Quadrupole-Quadrupole Metric., 2015. http://arxiv.org/abs/1507.04264

[27] F. Frutos-Alfaro. Approximate Kerr-like Metric with Quadrupole., 2015. http://arxiv.org/abs/1509.03698

[28] A. C. Hearn. *REDUCE* (User's and Contributed Packages Manual). Konrad-Zuse-Zentrum für Informationstechnik, Berlin, 1999. http://www.reduce-algebra.com/docs/reduce.pdf

[29] V. S. Manko and I. D. Novikov. Generalizations of the Kerr and Kerr-Newman Metrics Possessing an Arbitrary Set of Mass-Multipole Moments. Classical and Quantum Gravity, 9, 2477–2487, 1992. http://dx.doi.org/10.1088/0264-9381/9/11/013